

SYMBIO MYCOGRO TEE, FAIRWAY, PITCH AND LAWN FERTILISERS



Benefits

- ✓ *Fungi convert thatch, dead moss and grass clippings to humus*
- ✓ *Bacteria solubilise and retain nutrients*
- ✓ *Assists germination*
- ✓ *Creates the biology needed for perennial grass growth*

Part **organic fertilisers** with a unique blend of **rhizo bacteria and fungi**.

ORGANIC AND PART ORGANIC

- ✓ Symbio 12.2.9 50% Organic Slow Release

INORGANIC SLOW RELEASE

- ✓ Symbio 20.0.10 50% SCU Slow Release
- ✓ Symbio 15.2.15 20% SCU Slow Release
- ✓ Symbio 7.0.10 with additional ThatchEating fungi
- ✓ Symbio 10.0.10 with additional ThatchEating fungi



APPLICATION AND TIMING

Part organic fertilisers should be applied at 35gms -75 gms per sq.m. Inorganic fertilisers at 25gms – 35gm per sq.m and watered in in the absence of rain.

The fungi and bacteria are activated by moisture. For best results apply before rain is forecast or irrigate for 2-3 minutes after application.

These fertilisers may be applied via any standard fertiliser spreader

These fertilisers are designed to give immediate growth with a long lasting slow release element. Ideally applied in late spring or early summer, they can be applied during the summer months

12.2.9; 9.3.14; 15.2.15; 10.0.20

Hazard statements

Warning



Causes serious eye irritation.

Precautionary statements

Wash contaminated skin thoroughly after handling.

Wear protective gloves/protective clothing/eye protection/face protection.

Avoid breathing dust. In case of inadequate ventilation, wear respiratory protection

IF IN EYES: Rinse cautiously with water for several minutes.

Remove contact lenses, if present and easy to do. Continue rinsing. If eye irritation persists: Get medical advice/attention.

IF INHALED: If breathing is difficult, remove victim to fresh air and keep at rest in a position comfortable for breathing.

If experiencing respiratory symptoms call a doctor/physician.

12.2.9

Category 3 Material. Not for human consumption. Organic fertilisers or soil improvers: No grazing of farmed animals or use of crops as herbage during at least 21 days following application, in the case of pigs during at least 2 months following application.

PACK SIZE: 20kg • 25kg